

A Non Conventional Ergodic Theorem for a Nil-System

T. Ziegler

Abstract

We prove a non conventional pointwise convergence theorem for a nilsystem, and give an explicit formula for the limit.

1 Introduction

Non conventional averages of the form

$$\frac{1}{N} \sum_{n=1}^N \prod_{j=1}^k f_j(T^{jn}x) \quad (1)$$

were first introduced by Furstenberg in the ergodic theoretic proof of Szemerédi's theorem on arithmetic progressions [4]. Recent developments imply L^2 convergence of these averages to a limit for any k [6]. We are interested in pointwise convergence. The case $k = 1$ is the classical Birkhoff pointwise ergodic theorem. J. Bourgain has proved pointwise convergence for the case $k = 2$ [2]. Pointwise convergence is not known for $k \geq 3$.

Nilsystems arise in the study of the L^2 convergence of the averages (1). A nilspace is a homogeneous space of a nilpotent Lie group. A nilsystem consists of a finite measure nilspace N/Γ with a measure preserving transformation given by an element of N . It was shown by Furstenberg, Weiss [5], Conze, Lesigne [3] that 2-step nilsystems 'characterize' the L^2 behavior of the average (1) for the case $k = 3$. By 'characterize' we mean that the L^2 average can be calculated by projecting on an appropriate factor - the characteristic factor of the average in question. That k -step nil spaces play a role in the convergence of a $(k + 1)$ length multiple recurrence average can be expected, since for a k step nilspace N/Γ , $a^{k+1}x\Gamma$ is a function of $ax\Gamma, a^2x\Gamma, \dots, a^kx\Gamma$, and this provides a constraint on the $(k + 1)$ tuple $T^n x, \dots, T^{(k+1)}x$ in this case. For this reason it is particularly interesting to study these averages on nilsystems.

1.1 Statement of Result

We prove a pointwise convergence theorem for nilsystem: Let N be a k -step connected, simply connected nilpotent Lie group, Γ a discrete subgroup s.t. N/Γ is compact. Let $N^1 = N$, $N^i = [N^{i-1}, N]$ ($N^{k+1} = 1$), and set $\Gamma^i = \Gamma \cap N^i$. Then Γ^i is a discrete subgroup of N^i , and N^i/Γ^i is compact (Malcev [9]). Let m_i be the probability measure on N^i/Γ^i , invariant under translation by elements of N^i . Consider the measure preserving system $(N/\Gamma, m_1, T)$, where T denotes a translation by some element $a \in N$. In [8], E. Lesigne proved the following theorem for step 2 nilpotent groups:

Theorem 1.1 (*Lesigne*) *Suppose T acts ergodically, $f_1, f_2, f_3 \in L^\infty(N/\Gamma)$, then for almost all $x \in N$*

$$\begin{aligned} \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N T^n f_1(x\Gamma) T^{2n} f_2(x\Gamma) T^{3n} f_3(x\Gamma) \\ = \int_{N/\Gamma} \int_{N^2/\Gamma^2} f_1(xy_1\Gamma) f_2(xy_1^2 y_2\Gamma) f_3(xy_1^3 y_2^3\Gamma) dm_1(y_1\Gamma^1) dm_2(y_2\Gamma^2) \end{aligned}$$

We generalize this result and prove the following ergodic theorem:

Theorem 1.2 *Suppose T acts ergodically. If $f_1, \dots, f_{k+1} \in L^\infty(N/\Gamma)$, then for almost all $x \in N$*

$$\begin{aligned} \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \prod_{j=1}^{k+1} T^{jn} f_j(x\Gamma) \\ = \int_{N/\Gamma} \dots \int_{N^k/\Gamma^k} \prod_{j=1}^{k+1} f_j(x \prod_{i=1}^j y_i^{(j)} \Gamma) \prod_{j=1}^{k+1} dm_j(y_j\Gamma^j). \end{aligned}$$

The proof is a combination of the proof of Lesigne with the work of Leibman [7] on geometric sequences in groups. For each x we give the orbit of $(x, \dots, x)$ under $T \times \dots \times T^{k+1}$ the structure of a nilsystem, with the action of a transformation S_x . The main technical difficulty is with proving that the transformation S_x is ergodic a.e. x .

1.2 Acknowledgment

I thank David Lehari and Yoav Yaffe for helping me make the proofs more readable. This paper is a part of the author's PhD thesis. I thank my advisor Prof. Furstenberg for introducing me to ergodic theory, specifically

to questions concerning non conventional ergodic theorems, and for fruitful discussions.

2 Proof of Theorem 1.2

We define a set $\tilde{N}$ by $\tilde{N} = N^1 \times \dots \times N^k$. For $(x_1, \dots, x_k), (y_1, \dots, y_k) \in \tilde{N}$ we define

$$z_1 = x_1 y_1, \\ \prod_{j=1}^i z_j^{(i)} = \prod_{j=1}^i x_j^{(i)} \prod_{j=1}^i y_j^{(i)}.$$

$$(x_1, \dots, x_k) \star (y_1, \dots, y_k) = (z_1, \dots, z_k),$$

One should think of $(x_1, \dots, x_k), (y_1, \dots, y_k)$ as representing the vectors

$$(x_1, x_1^2 x_2, \dots, \prod_{j=1}^k x_j^{(k)}), \quad (y_1, y_1^2 y_2, \dots, \prod_{j=1}^k y_j^{(k)})$$

respectively. Then $(z_1, \dots, z_k)$ represents the coordinate product of these vectors. By Leibman [7], $\tilde{N}$ is a group under this multiplication, thus $\tilde{N}$ is a nilpotent connected simply connected Lie group. Let $\tilde{\Gamma} = \Gamma^1 \times \dots \times \Gamma^k$, then $(\tilde{\Gamma}, \star)$ is a discrete subgroup of $\tilde{N}$, and the quotient $\tilde{N}/\tilde{\Gamma}$ is compact. For $x \in N$ we define a transformation $S_x : \tilde{N}/\tilde{\Gamma} \rightarrow \tilde{N}/\tilde{\Gamma}$ by:

$$S_x((y_1, \dots, y_k) \star \tilde{\Gamma}) = ((a[a, x], e, \dots, e) \star (y_1, \dots, y_k)) \star \tilde{\Gamma},$$

and a function $I : \tilde{N} \rightarrow (N/\Gamma)^{k+1}$ by:

$$I((y_1, \dots, y_k)) = (xy_1\Gamma, xy_1^2 y_2\Gamma, \dots, x \prod_{j=1}^k y_j^{(k+1)} \Gamma).$$

Let $(\gamma_1, \dots, \gamma_k) \in \tilde{\Gamma}$ then we have for all $(y_1, \dots, y_k) \in \tilde{N}$:

$$\begin{aligned} & I((y_1, \dots, y_k) \star (\gamma_1, \dots, \gamma_k)) \\ &= (xy_1\gamma_1\Gamma, xy_1^2 y_2\gamma_1^2\gamma_2\Gamma, \dots, x \prod_{j=1}^k y_j^{(k+1)} \prod_{j=1}^k \gamma_j^{(k+1)} \Gamma) \\ &= (xy_1\Gamma, xy_1^2 y_2\Gamma, \dots, x \prod_{j=1}^k y_j^{(k+1)} \Gamma) \\ &= I((y_1, \dots, y_k)), \end{aligned}$$

thus I can be defined on $\tilde{N}/\tilde{\Gamma}$. Let $\tilde{m}$ denote the probability measure on $\tilde{N}/\tilde{\Gamma}$, invariant under the action of $\tilde{N}$, then $\tilde{m} = m_1 \times \dots \times m_k$. Denote by m_x, A_x the images of $\tilde{m}, \tilde{N}/\tilde{\Gamma}$ under I . The function I is an isomorphism of the systems $(A_x, m_x, T \times T^2 \times \dots \times T^{k+1})$ and $(\tilde{N}/\tilde{\Gamma}, \tilde{m}, S_x)$ since

$$\begin{aligned}
& I(S_x((y_1, \dots, y_k) \star \tilde{\Gamma})) \\
&= I((a[a, x], e, \dots, e) \star (y_1, \dots, y_k) \star \tilde{\Gamma}) \\
&= (xa[a, x]y_1\Gamma, x(a[a, x])^2y_1^2y_2\Gamma, \dots, x(a[a, x])^{k+1} \prod_{j=1}^k y_j^{\binom{k+1}{j}} \Gamma) \\
&= (axy_1\Gamma, a^2xy_1^2y_2\Gamma, \dots, a^{k+1}x \prod_{j=1}^k y_j^{\binom{k+1}{j}} \Gamma) \\
&= (T \times T^2 \times \dots \times T^{k+1})(I((y_1, \dots, y_k) \star \tilde{\Gamma})).
\end{aligned}$$

If the action of S_x is ergodic, then by Parry [10], [11] the m.p.s $(\tilde{N}/\tilde{\Gamma}, \tilde{m}, S_x)$ is uniquely ergodic, and thus the m.p.s $(A_x, m_x, T \times T^2 \times \dots \times T^{k+1})$ is uniquely ergodic. For each such x we have: for all $(y_1, \dots, y_k) \in \tilde{N}$, for all continuous functions F on $(N/\Gamma)^{k+1}$,

$$\begin{aligned}
& \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N F(a^n xy_1\Gamma, a^{2n} xy_1^2 y_2\Gamma, \dots, a^{(k+1)n} x \prod_{j=1}^k y_j^{\binom{k+1}{j}} \Gamma) \\
&= \int_{N/\Gamma} \dots \int_{N_k/\Gamma_k} F(xy_1\Gamma, xy_1^2 y_2\Gamma, \dots, x \prod_{j=1}^k y_j^{\binom{k+1}{j}} \Gamma) \prod_{j=1}^k dm_j(y_j\Gamma_j).
\end{aligned}$$

In particular, for all such F

$$\begin{aligned}
& \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N F(a^n x\Gamma, a^{2n} x\Gamma, \dots, a^{(k+1)n} x\Gamma) \\
&= \int_{N/\Gamma} \dots \int_{N_k/\Gamma_k} F(xy_1\Gamma, xy_1^2 y_2\Gamma, \dots, x \prod_{j=1}^k y_j^{\binom{k+1}{j}} \Gamma) \\
&\quad dm_1(y_1\Gamma) \dots dm_k(y_k\Gamma).
\end{aligned}$$

If S_x is ergodic then we proved the theorem for $f_1, \dots, f_k$ continuous. For passing to bounded measurable functions see [3] lemma 3 p. 173.

In order to finish the proof of the theorem, we must show that the action of S_x is ergodic for a.a. $x \in N$.

2.1 Ergodicity of S_x

We wish to show that the action of S_x on $\tilde{N}$ is ergodic for almost all (Haar) $x \in N$. Let σ be a character of $\tilde{N}$ s.t. σ vanishes on $\tilde{N}' \star \tilde{\Gamma}$. By Green [1] we must show that if $\sigma \not\equiv 1$ then

$$m_h\{x \in N : \sigma((a[a, x], e, \dots, e)) = 1\} = 0$$

(as there is only countable number of characters on $\tilde{N}/\tilde{N}'\tilde{\Gamma}$). Suppose this set is of positive measure. Let $\psi(x) = \sigma((a[a, x], e, \dots, e))$, then ψ is an analytic function on N , therefore $\psi(x) = 1$ on a set of positive measure implies $\psi \equiv 1$. We define the functions σ_i on N^i by

$$\sigma_i(y_i) = \sigma((e, \dots, e, y_i, e, \dots, e)).$$

We have

$$\sigma((y_1, \dots, y_k)) = \prod_{i=1}^k \sigma_i(y_i).$$

For $y_1, \dots, y_n \in N$ we define $R(y_1, \dots, y_n)$ as follows:

$$\begin{aligned} R^0(y_1, \dots, y_n) &= \{y_1, \dots, y_n\} \\ R^j(y_1, \dots, y_n) &= \bigcup_{i,l=0}^{j-1} \{[R^i, R^l]\} \\ R(y_1, \dots, y_n) &= \bigcup_{i=1}^k R^i(y_1, \dots, y_n). \end{aligned}$$

In other words, $R(y_1, \dots, y_n)$ is the set of commutators involving $y_1, \dots, y_n$. For all x

$$\begin{aligned} 1 &= \psi(x) = \sigma((a[a, x], e, \dots, e)) = \sigma_1(a[a, x]) \\ &= \sigma_1(a)\sigma_1([a, x])\sigma_2(c_2(x)) \dots \sigma_k(c_k(x)), \end{aligned}$$

where $c_i(x)$ are products of elements of $R(a, [a, x])$.

Remark 2.1 $\sigma_i(xy) = \sigma_i(x)\sigma_i(y)$ modulo ‘corrections’ in σ_j , $j > i$ computed on commutators involving x, y .

Setting $x = e$ we get $\sigma(a) = 1$, thus

$$\begin{aligned}\sigma_1([a, x]) &= \sigma_2(c_2(x))^{-1} \dots \sigma_k^{-1}(c_k(x)) \\ &= \prod_{i=2}^k \prod_{m=1}^{l_i} \sigma_i(r_{im}(x)),\end{aligned}\tag{2}$$

for all $x \in N$, where $r_{im}(x) \in R(a, [a, x])$ (as $R(a, [a, x])$ is closed under $[,]$). We will show that σ_i are all trivial, and thus σ is trivial.

Notation 1 We use $[x_1, \dots, x_n]$ for $[\dots [[x_1, x_2], x_3], x_4], \dots, x_n]$.

Notation 2 Let $\mathcal{S} = \{(s_1, \dots, s_j) : 0 \leq j \leq k, 1 \leq s_n \leq k\}$. For $A = (s_1, \dots, s_n) \in \mathcal{S}$ define C_A by:

$$\begin{aligned}C_A^0 &= a \\ C_A^j &= \{[C_A^{j-1}, x_j]^{\pm 1} : x_j \in N^{s_j}\} \\ C_A &= C_A^n\end{aligned}$$

Example : If $A = (1, 3)$ then $C_A = \{[[a, x]^{\pm 1}, y]^{\pm 1}\}$ where $x \in N, y \in N^3$.

Notation 3 1. For $A = (s_1, \dots, s_j) \in \mathcal{S}$, $|A| = \sum s_j$

2. For $A \in \mathcal{S}$, $(i : A)$ means: $\forall v \in (C_A \cap N^i) : \sigma_i(v) = 1$ (if $|A| < i - 1$ the intersection is empty).

3. $j^{\geq} (>)$ is some $l \geq j$ ($l > j$).

4. Let X be either $j, j^{\geq}, j^{>}$, then $X+$ is a sequence of length ≥ 1 , whose entries are of type $j, j^{\geq}, j^{>}$ respectively.

5. For $x \in N^j$, $|x| = j$.

Proposition 2.2 $\forall A \in \mathcal{S}, 1 \leq i \leq k : (i : A)$.

Proof: We will need the following lemma, which is based on the fact that σ is multiplicative and thus $\sigma \equiv 1$ on $\tilde{N}'$:

Lemma 2.3 Let $x \in N, y \in N^n$. We define

$$H(x, y) = \bigcup_{i=2}^k R^i(y_1, \dots, y_n) \quad (= R(x, y) \setminus \{[x, y]^{\pm 1}\}).$$

If $\sigma_i(h) = 1 \ \forall i > 1, h \in H(x, y) \cap N^i$, then $\sigma_m([x, y])^m = \sigma_{m+1}([x, y])^{m+1}$, for all $m \leq n$.

Proof: We denote by $\tilde{H}(x, y)$ the set of finite products of elements of $H(x, y)$. By remark (2.1) and since $H(x, y)$ is closed under commutators, $\sigma_i(\tilde{h}) = 1$ ($i > 1$) for all $\tilde{h} \in \tilde{H}(x, y) \cap N^i$. As $\sigma|_{\tilde{N}'} \equiv 1$, we have

$$\begin{aligned} 1 &= \sigma((x^{-1}, e, \dots, e) \star (e, \dots, y^{-1}, \dots, e) \star (x, e, \dots, e) \star (e, \dots, y^m, \dots, e)) \\ &= \sigma(z_1, \dots, z_k) = \prod_{i=1}^k \sigma_i(z_i), \end{aligned}$$

we calculate the z_i :

$$\begin{aligned} z_1 &= \dots = z_{m-1} = e, \\ z_m &= [x^m, y] = [x, y]^m \tilde{h}_0 \end{aligned}$$

for some $\tilde{h}_0 \in \tilde{H}(x, y)$.

$$\begin{aligned} z_m^{(m+1)} z_{m+1} &= [x^{(m+1)}, y^{(m+1)}] = [x, y]^{(m+1)^2} \tilde{h}_1' \Rightarrow \\ z_{m+1} &= [x, y]^{(m+1)} \tilde{h}_1 \end{aligned}$$

for some $\tilde{h}_1, \tilde{h}_1' \in \tilde{H}(x, y)$. We show by induction that $z_{m+j} = \tilde{h}_j$ for $j > 1$, $\tilde{h}_j \in \tilde{H}(x, y)$. For $j = 2$:

$$\begin{aligned} z_m^{\binom{m+2}{2}} z_{m+1}^{(m+2)} z_{m+2} &= [x^{(m+2)}, y^{\binom{m+2}{2}}] = [x, y]^{(m+1)\binom{m+2}{2}} \tilde{h}_2' \Rightarrow \\ z_{m+2} &= [x, y]^{-m\binom{m+2}{2} - (m+1)(m+2) + (m+2)\binom{m+2}{2}} \tilde{h}_2 = \tilde{h}_2 \end{aligned}$$

for some $\tilde{h}_2, \tilde{h}_2' \in \tilde{H}(x, y)$. Suppose it is true for $2 \leq i < j$, we have

$$\begin{aligned} z_m^{\binom{m+j}{j}} z_{m+1}^{\binom{m+j}{j-1}} \dots z_{m+j} &= [x^{(m+j)}, y^{\binom{m+j}{j}}] = [x, y]^{(m+j)\binom{m+j}{j}} \tilde{h}_j' \Rightarrow \\ z_{m+j} &= [x, y]^{-m\binom{m+j}{j} - (m+1)\binom{m+j}{j-1} + (m+j)\binom{m+j}{j}} \tilde{h}_j = \tilde{h}_j \end{aligned}$$

for some $\tilde{h}_j, \tilde{h}_j' \in \tilde{H}(x, y)$. Thus

$$\begin{aligned} 1 &= \prod_{i=1}^k \sigma_i(z_i) = \sigma_m([x, y]^m \tilde{h}_0) \sigma_{m+1}([x, y]^{m+1} \tilde{h}_1) \prod_{i=m+2}^k \sigma_i(\tilde{h}_{i-m}) \\ &= \sigma_m([x, y]^m) \sigma_{m+1}([x, y]^{m+1}), \end{aligned}$$

as $\tilde{H}([x, y], \tilde{h}_0), \tilde{H}([x, y], \tilde{h}_1) \subset \tilde{H}(x, y)$. ♠

Corollary 2.4 *If $\sigma_1([x, y]) = 1$, and $\sigma_i(h) = 1$ for all $h \in H(x, y) \cap N^i$ then $\sigma_i([x, y]) = 1$ for all i s.t. $[x, y] \in N^i$.*

Proof: By lemma 2.3, $\sigma_i([x, y])^i = 1$. N is connected, and $\sigma_i([e, y]) = 1$. ♠

Proposition 2.5 *We prove the following rules:*

1. $|A| \geq k \implies \forall i : (i : A)$. (*triviality rule*)
2. $\forall i : (i : j^{>+}) \implies (1 : j)$. (*reducing rule*)
3. $(1 : (A, l)) \ \& \ j > l \ \& \ \forall i : (i : (A, l, j^{\geq+})) \ \& \ \forall i : (i : (A, j - 1, j^{\geq+})) \implies (1 : (A, l, j - 1)) \ \& \ (1 : (A, j - 1, l))$. (*expanding rule*)
4. $(1 : (A, j)) \ \& \ \forall i : (i : (A, j^{\geq}, j^{\geq+})) \ \& \ \forall i : (i : (A, j^{>})) \implies \forall i : (i : (A, j))$.

Proof: Let $A = (s_1, \dots, s_j)$, $0 \leq j \leq k$, $x_n \in N^{s_n}$.

1. $|A| \geq k$ implies that $[a, x_1, \dots, x_j] = 1$ thus $\sigma_i([a, x_1, \dots, x_j]) = 1$.
2. Let $x \in N^j$.

We use equation 2:

$$\sigma_1([a, x]) = \prod_{i=2}^k \prod_{m=1}^{l_i} \sigma_i(r_{i_m}(x)),$$

where $r_{i_m}(x) \in R(a, [a, x])$. Denote $b = [a, x]$, then $r_{i_m}(x) \in R(a, b)$. Looking at the a appearing in a minimal number of brackets, $r_{i_m}(x)$ is of the form: $[a, y_1, \dots, y_m]$ where $m \geq 1$, $|y_n| \geq |b| > |x|$, and is thus of type $(j^{>+})$.

3. Let $v = [a, x_1, \dots, x_j]$, $y \in N^l$, $z \in N^{j-1}$ then $yz \in N^l$.

$$\begin{aligned} 1 &= \sigma_1([v, yz]) = \sigma_1([v, y][v, y, z][v, z]) \\ &= \sigma_1([v, y])\sigma_1([v, y, z])\sigma_1([v, z]) \prod_{i=2}^k \prod_{m=1}^{l_i} \sigma_i(b_{i_m}(y, z, v)) \\ &= \sigma_1([v, y, z]) \prod_{i=2}^k \prod_{m=1}^{l_i} \sigma_i(b_{i_m}(y, z, v)), \end{aligned}$$

where $b_{i_m} \in R([v, y], [v, z], [v, y, z])$. If $[v, y]$ appears in $b_{i_m}(y, z, v)$ then w.l.o.g.

$$b_{i_m}(y, z, v) = [v, y, y_1, \dots, y_m],$$

where $m \geq 1$, $|y_n| \geq j$ (look at the $[v, y]$ appearing in the minimal number of brackets) This is of type $(A, l, j^{\geq +})$. Otherwise $[v, y]$ does not appear in $b_{i_m}(y, z, v)$, thus w.l.o.g.


$$b_{i_m}(y, z, v) = [v, z, z_1, \dots, z_m],$$

where $m \geq 1$, $|z_n| \geq j$. This is of type $(A, j-1, j^{\geq +})$. For $(A, j-1, l)$, do the same with $[v, zy]$.

4. Let v be of type A , $x \in N^j$. By lemma 2.3, we must show that for $h \in H(v, x) \cap N^i$, $\sigma_i(h) = 1$. But any such h is either of the form $[v, r(v, x)]$ where $r(v, x) \in R(v, x)$ and thus of type $(A, j^>)$, or $[v, z_1, \dots, z_m]$, $m \geq 1$, $|z_n| \geq j$, and thus of type $(A, j^{\geq}, j^{\geq +})$.

Proposition 2.6 $(1 : (A, l)) \implies (i : (A, l^{\geq +}))$, where $|A| \geq 0$, $l \geq 1$.

Proof: We prove this by induction on $|A|$, l . For $|A| = k$ any l it is clear (as the commutator is trivial). Suppose the statement is true for $|A| > d$, any l . For $|A| = d$, $l = k - d$ it is clear (as the commutator is trivial). Suppose the statement is true for $|A| = d$, any $j > l$. We follow the following scheme:



Explanation for numbered arrows::

1. $k - |(A, l)| < l$ implies $(i : (A, l^{\geq}, l^{\geq+}))$ by the triviality rule 1. $(1 : (A, l))$ implies $(1 : (A, j))$ for $j > l$ (as $x \in N^j \Rightarrow x \in N^l$) By the induction hypothesis $(i : (A, j))$ for $j > l$, and by rule 4 $(i : (A, l))$.
2. By expanding rule 3 and triviality rule 1.
3. By the induction hypothesis (as $|A, j| > |A|$).
4. By the induction hypothesis (as $|A, l| > |A|$).
5. By expanding rule: $(j - 1) \geq l$, $(1 : (A, l))$, $(i : (A, l, j^{\geq+}))$, and $(i : (A, (j - 1), j^{\geq+}))$ as if $j - 1 = l$ it is the previous condition, and if $j - 1 > l$ then $(1 : (A, l))$ implies $(1 : (A, j - 1))$ which implies by the induction $(i : (A, j - 1^{\geq+}))$.
6. Repeat the procedure until $j = l$.
7. By this procedure we now know $(i : A, l^{\geq}, l^{\geq+})$. Repeat the argument in 1.

Proof of proposition 2.2 We prove this by induction on j : $(1 : k)$ by triviality rule 1. $(1 : j) \Rightarrow \forall i(i : j^{\geq+})$ by proposition 2.6 ($|A| = 0$). By rule 2 this implies $(1 : j - 1)$. ♠

Proposition 2.7 *For all i , $\sigma_i \equiv 1$.*

Proof: We prove this by induction on k . By proposition 2.2, we have

$$\sigma_k([x_1, \dots, \overset{m}{a}, \dots, x_{k-1}]) = 1 \quad \forall x_1, \dots, x_{k-1} \in N, \quad 1 \leq m \leq k \quad (3)$$

Since $\sigma|_{\Gamma} \equiv 1$,

$$\sigma_k([\gamma_1, \dots, \gamma_k]) = 1 \quad \forall \gamma_1, \dots, \gamma_k \in \Gamma.$$

From (3)

$$\sigma_k([a, \gamma_2, \dots, \gamma_k]) = 1 \quad \forall \gamma_2, \dots, \gamma_k \in \Gamma.$$

Since σ_k is a character on N^k , and a acts ergodically on N/Γ , we have

$$\sigma_k([x_1, \gamma_2, \dots, \gamma_k]) = 1 \quad \forall \gamma_2, \dots, \gamma_k \in \Gamma, \quad x_1 \in N.$$

Using (3) and induction, we get

$$\sigma_k([x_1, \dots, x_k]) = 1 \quad \forall x_1, \dots, x_k \in N,$$

thus $\sigma_k \equiv 1$. Now $\sigma_k, \dots, \sigma_{j+1} \equiv 1$ implies that σ_j is a character. For $j > 1$, by Proposition 2.2, we have

$$\sigma_j([x_1, \dots, \overset{m}{a}, \dots, x_{j-1}]) = 1 \quad \forall x_1, \dots, x_{j-1} \in N, \quad 1 \leq m \leq j,$$

and by the same argument as above we have $\sigma_j \equiv 1$. For $j = 1$, σ_1 is a character which satisfies

$$\sigma_1(a) = 1, \quad \sigma_1|_{\Gamma} \equiv 1,$$

thus $\sigma_1 \equiv 1$. ♠

References

- [1] Auslander, L., Green, L., Hahn, F. *Flows on homogeneous spaces*. Annals of Mathematics Studies, No. 53 (1963).
- [2] Bourgain, J. *Pointwise ergodic theorems for arithmetic sets*. Inst. Hautes études Sci. Publ. Math. No. 69: 5-45 (1989).
- [3] Conze, J.P., Lesigne, E. *Théorèmes ergodiques pour des mesures diagonales*. Bull. Soc. Math. France 112 no. 2: 143-175 (1984).
- [4] Furstenberg, H. *Ergodic behavior of diagonal measures and a theorem of Szemerédi on arithmetic progressions*. J. Analyse Math. 31: 204-256 (1977).
- [5] Furstenberg, H., Weiss B. *A mean ergodic theorem for $(1/N) \sum_{n=1}^N f(T^n x)g(T^{n^2} x)$* . Convergence in ergodic theory and probability Ohio State Univ. Math. Res. Inst. Publ., 5, de Gruyter, Berlin: 193-227 (1996).
- [6] Host, B. , Kra, B : personal communication.
- [7] Leibman, A *Polynomial sequences in groups*. Journal of Algebra, 201: 189-206 (1998).
- [8] Lesigne, E. *Theorems ergodiques pour une translation sur une nilvariété* Erg. Thm and Dyn.Sys, 9(1): 115-126 (1989).

- [9] Malcev, A.I. *On a class of homogeneous spaces*. Amer. Math. Soc. Translation no. 39 (1951).
- [10] Parry, W. (1969) *Ergodic properties of affine transformations and flows on nilmanifolds*. Amer. J. Math. 91 :757-771.
- [11] Parry, W. *Dynamical systems on nilmanifolds*. Bull. London Math. Soc. 2: 37-40 (1970).